arXiv:math/9812062v1 [math.FA] 9 Dec 1998

Injective isometries in Orlicz spaces

Beata Randrianantoanina

Abstract. We show that injective isometries in Orlicz space L_M have to preserve disjointness, provided that Orlicz function M satisfies Δ_2-condition, has a continuous second derivative M'', satisfies another "smoothness type" condition and either $\lim_{t\to 0} M''(t) = \infty$ or $M''(0) = 0$ and $M''(t) > 0$ for all $t > 0$. The fact that surjective isometries of any rearrangement-invariant function space have to preserve disjointness has been determined before. However dropping the assumption of surjectivity invalidates the general method. In this paper we use a differential technique.

1. Introduction

The study of isometries of Banach spaces goes back to Banach's 1932 treatise on linear operators [**Ban**] and since then it received much attention in the literature, see the survey [**FJ**] with its over 300 references. We just mention here the results most closely related to the present work.

Banach showed that in separable $L_p(\Omega, \mu), p \neq 2$, any isometry $T : L_p(\Omega, \mu) \longrightarrow L_p(\Omega, \mu)$ is of the form

$$Tf(\omega) = h(\omega)f(\sigma(\omega))$$

where $\sigma : \Omega \longrightarrow \Omega$ is a Borel automorphism of Ω and h is a scalar function on Ω. Banach obtained this result by showing that every isometry T of L_p has to preserve disjointness i.e. if $f, g \in L_p$ are such that $\mu(\operatorname{supp} f \cap \operatorname{supp} g) = 0$ then also $\mu(\operatorname{supp} Tf \cap \operatorname{supp} Tg) = 0$ (cf. also [**Lam**]). To achieve this he characterized disjointness of functions f, g through the differential properties of the function $N(\alpha) = \|f + \alpha g\|$. Similar technique was later applied by Koldobsky [**Kol**] to study injective isometries of $L_p(L_q)$ and Kamińska [**Kam**], who observed that isometries of Orlicz spaces L_M preserve disjointness under assumptions that both M and M' are strictly convex, $M'(0) = M''(0) = 0$ and M satisfies Δ_2-condition.

In the present paper we adapt Banach's differential technique to Orlicz spaces L_M, where M satisfies Δ_2-condition, another "smoothness type" condition (see Definition 2.2) and has a continuous second derivative M''. We note that unlike Banach and Koldobsky we do not obtain conditions which are equivalent to the disjointness of supports of functions f, g. In the case when $M''(0) = 0$ we only describe some conditions which are necessary for disjointness and some conditions which are sufficient for disjointness, which together enable us to characterize the

---

1991 *Mathematics Subject Classification.* 46E30,46B04.

©0000 (copyright holder)

injective isometries. The argument in the case $\lim_{t\to 0} M''(t) = \infty$ is even more delicate and depends on the sum of sufficient conditions and necessary conditions for containment of supports of f and g. The lack of isometric conditions equivalent to disjointness should not be really surprising in view of the classical Bohnenblust's characterization of L_p-spaces [**LT**, Theorem 1.b.7].

We conclude the introduction by recalling that surjective isometries of complex Orlicz spaces were described by Lumer [**L1, L2**] (reflexive case) and Zaidenberg [**Z**]. Jamison, Kamińska and P.K. Lin [**JKL**] studied surjective isometries of complex Musielak-Orlicz spaces and real Nakano spaces. The form of surjective isometries of real Orlicz spaces follows from the description of surjective isometries of real rearrangement-invariant spaces [**KR**].

The results of present paper are valid in both complex and real case.

ACKNOWLEDGMENTS . I wish to thank Professor A. Koldobsky for introducing me to this problem and for suggesting the use of the differential technique. I am also grateful to Professors Y. Abramovich and A. Kamińska for valuable discussions, and to Professor D. Burke for help with drawing Figures 1 and 2.

Part of this work was completed while the author participated in the Workshop in Linear Analysis and Probability at Texas A&M University, College Station, Texas in the Summer 1997 organized by Professors W. Johnson, D. Larson, G. Pisier and J. Zinn. I wish to thank the organizers for their hospitality and support.

2. Definitions and preliminary lemmas

We follow standard definitions and notations as may be found e.g. in [**KrR**] or [**Ch**]. We recall the basic definitions below.

We say that a function $M : [0,\infty) \longrightarrow [0,\infty)$ is an *Orlicz function* if M is convex, $M(0) = 0$, $M(1) = 1$, $\lim_{u\to 0} M(u)/u = 0$ and $\lim_{u\to\infty} M(u)/u = \infty$.

The Orlicz function M generates the *Luxemburg norm* defined for scalar valued functions on Ω by:

$$\|f\|_M = \inf\{\lambda : \int_\Omega M\left(\frac{|f(u)|}{\lambda}\right) d\mu \le 1\}.$$

The *Orlicz space* L_M is the space of (equivalence classes of) measurable functions f with $\|f\|_M < \infty$.

We say that two Orlicz functions M_1 and M_2 are *equivalent* if there exist $u_0 > 0, k, l > 0$ such that for all $u > u_0$

$$M_2(ku) \le M_1(u) \le M(lu).$$

This condition is of importance since Orlicz spaces L_{M_1}, L_{M_2} are isomorphic if and only if the Orlicz functions M_1, M_2 are equivalent.

It is well known (see e.g. [**Ch**]) that any Orlicz function M can be "smoothed out", that is for any M there exists an equivalent Orlicz function M_1 such that M_1 is twice differentiable and $M_1''(u) > 0$ for all $u > 0$. Moreover, given any $\varepsilon > 0$ it is possible to choose M_1 so that L_M and L_{M_1} are $(1+\varepsilon)-$isomorphic to each other [**Ch**].

We say that the Orlicz function $M(u)$ *satisfies the* Δ_2 *condition for large values of* u if there exist constants $k > 0$ and $u_0 \ge 0$ such that for all $u \ge u_0$

$$M(2u) \le kM(u).$$

If the Orlicz function M satisfies the Δ_2 condition then every Orlicz function M_1 equivalent to M also satisfies the Δ_2 condition.

Note that Orlicz function M is convex so it has the right derivative M'. Krasnoselskii and Rutickii provide the following characterization of the Δ_2-condition in terms of M'.

PROPOSITION 2.1. [**KrR**, Theorem 4.1] *A necessary and sufficient condition that the Orlicz function $M(u)$ satisfy the Δ_2-condition is that there exist constants α and $u_0 \geq 0$ such that, for $u \geq u_0$*

$$\frac{uM'(u)}{M(u)} < \alpha, \tag{2.1}$$

where M' denotes the right derivative of M.

Moreover, if (2.1) *is satisfied then $M(2u) \leq 2^\alpha M(u)$ for $u \geq u_0$.*

Functions which satisfy the Δ_2-condition do not increase more rapidly than polynomials [**KrR**, p. 24]. In fact there exists a constant C so that for all $u \geq u_0$

$$M(u) < Cu^\alpha, \tag{2.2}$$

where α and u_0 are the same as in (2.1).

It is also true (see [**Ch**] or [**KrR**]) that for any Orlicz function M and for any $u > 0$

$$\frac{uM'(u)}{M(u)} > 1, \tag{2.3}$$

We now introduce another condition which on one hand is very similar to (2.1), but on the other hand is in its nature of "smoothness type", as we explain below.

DEFINITION 2.2. Assume that the Orlicz function M is twice differentiable and that M satisfies the Δ_2- condition. We say that M *satisfies condition* Δ_{2+} if there exist constants $\beta > 0$ and $u_0 \geq 0$ such that for all $u \geq u_0$

$$\frac{uM''(u)}{M'(u)} < \beta. \tag{2.4}$$

Condition Δ_{2+} is very important for us because we can prove our isometry results only for Orlicz spaces L_M, such that M satisfies Δ_{2+}. Thus we wish to observe that this condition is of "smoothness type" in the following sense:

(i) for every function M which satisfies condition Δ_2 there exists an equivalent Orlicz function M_1 which does satisfy Δ_{2+}, see Lemma 2.3; However, we do not know whether for every $\varepsilon > 0$ it is possible to choose M_1 so that it is $(1+\varepsilon)-$equivalent with M,

(ii) for every Orlicz function M which satisfies Δ_{2+} there exists an equivalent (even up to an arbitrary $\varepsilon > 0$) Orlicz function M_1 which does not satisfy Δ_{2+} (see Lemma 2.5).

LEMMA 2.3. *Let M be any Orlicz function which satisfies condition Δ_2. Then there exists an Orlicz function M_1 which is equivalent to M and satisfies condition Δ_{2+}.*

REMARK 2.4. We do not know whether for every $\varepsilon > 0$ and every M satisfying Δ_2 it is possible to choose M_1 satisfying Δ_{2+} and $(1+\varepsilon)-$equivalent to M.

Proof. Define M_1' by (see Figure 1):

$$M_1'(u) = \begin{cases} M'(u) & \text{if } u \le 2 \\ f_k(u) & \text{if } 2^k \le u \le 2^k+1 \ , k \in \mathbb{N} \\ M'(2^k) + (u-2^k)\frac{M'(2^{k+1})-M'(2^k)}{2^k} & \text{if } 2^k+1 < u < 2^{k+1} \ , k \in \mathbb{N} \end{cases}$$

where $f_k : \mathbb{R} \longrightarrow \mathbb{R}$ is an increasing, continuous, differentiable function such that f_k' is monotone on $[2^k, 2^k+1]$ and

$$\begin{aligned} f_k(2^k) &= M'(2^k) \\ f_k(2^k+1) &= M'(2^k) + \frac{M'(2^{k+1}) - M'(2^k)}{2^k} \\ f_k'(2^k) &= \frac{M'(2^k) - M'(2^{k-1})}{2^{k-1}} \\ f_k'(2^k+1) &= \frac{M'(2^{k+1}) - M'(2^k)}{2^k}. \end{aligned}$$



Figure 1.

Thus $M_1'(u)$ is increasing, continuous and differentiable for all $u \in \mathbb{R}_+$. Then

$$M_1(u) = \int_0^u M_1'(t)\ dt.$$

We first check that M_1 is equivalent to M. Indeed, for any $u \ge 2$ let $k \in \mathbb{N}$ be such that $2^k \le u < 2^{k+1}$. Then we have:

$$\begin{aligned} M_1(u) = \int_0^u M_1'(t)\ dt &\ge \int_0^2 M'(t)\ dt + \sum_{j=1}^{k} 2^j M'(2^j) \\ &\text{by (2.3)} \\ &> \sum_{j=1}^{k} M(2^j) > M(2^k) \ge M(\frac{u}{2}) \end{aligned}$$

Also we have:

$$\begin{aligned}
M_1(u) &= \int_0^u M_1'(t)\ dt \\
&\leq \int_0^2 M'(t)\ dt + \sum_{j=1}^{k} 2^j M'(2^{j+1}) + (u-2^k)M'(2^{k+1}) \\
&\text{by (2.1)} \\
&\leq M(1) + \frac{\alpha}{2}\sum_{j=1}^{k+1} M(2^{j+1}) \leq \frac{1}{2} + \frac{\alpha}{2}\sum_{l=1}^{k+2} M(2^l) \\
&\text{by convexity of } M \\
&\leq \frac{1}{2} + \frac{\alpha}{2}\sum_{l=1}^{k+2} \frac{M(2^{k+2})}{2^{k+2-l}} \\
&\leq \frac{1}{2} + \frac{\alpha}{2}M(2^{k+2}) \leq \frac{1}{2} + \frac{\alpha}{2}M(4u) < \alpha M(4u) \\
&\text{since } \alpha > 1 \\
&\leq M(4\alpha u)
\end{aligned}$$

Since for $u \leq 2$ $M(u) = M_1(u)$, we conclude that for all $u \geq 0$

$$M(\frac{u}{2}) \leq M_1(u) \leq M(4\alpha u)$$

i.e. M_1 is equivalent to M.

Next we show that M_1 satisfies condition Δ_{2+}.

Let $u \geq 2$ and $k \in \mathbb{N}$ be such that $2^k \leq u < 2^{k+1}$. Then, if $2^k + 1 \leq u$ we have:

$$\begin{aligned}
\frac{uM_1''(u)}{M_1'(u)} &\leq \frac{2^{k+1}\frac{M'(2^{k+1})-M'(2^k)}{2^k}}{M'(2^k)} = \frac{2M'(2^{k+1})}{M'(2^k)} - 2 \\
&\text{by (2.1) and (2.3)} \\
&\leq \frac{2\frac{\alpha M(2^{k+1})}{2^{k+1}}}{\frac{M(2^k)}{2^k}} = \frac{\alpha M(2^{k+1})}{M(2^k)} \\
&\text{by Proposition 2.1} \\
&\leq \alpha 2^{\alpha}.
\end{aligned}\tag{2.5}$$

If $2^k \leq u \leq 2^k + 1$, then $M_1'(u) = f_k(u)$ and we consider two cases:

(1) If f_k' is increasing on $[2^k, 2^k+1]$ then

$$f_k'(u) \leq f_k'(2^k+1) = \frac{M'(2^{k+1}) - M'(2^k)}{2^k}$$

and

$$\begin{aligned}
\frac{uM_1''(u)}{M_1'(u)} = \frac{uf_k'(u)}{f_k(u)} &\leq \frac{2^k + 1\frac{M'(2^{k+1})-M'(2^k)}{2^k}}{M'(2^k)} \\
&\text{by (2.5)} \\
&\leq \alpha 2^{\alpha}.
\end{aligned}$$

(2) If f_k' is decreasing on $[2^k, 2^k+1]$ then

$$f_k'(u) \leq f_k'(2^k) = \frac{M'(2^k) - M'(2^{k-1})}{2^{k-1}}$$

and

$$\begin{aligned} \frac{uM_1''(u)}{M_1'(u)} = \frac{uf_k'(u)}{f_k(u)} &\leq \frac{(2^k+1)\frac{M'(2^k)-M'(2^{k-1})}{2^{k-1}}}{M'(2^k)} \\ &\leq \frac{2\cdot 2^k \frac{M'(2^k)-M'(2^{k-1})}{2^{k-1}}}{M'(2^{k-1})} \\ &\text{by (2.5) applied to } (k-1) \\ &\leq 2\alpha 2^\alpha. \end{aligned}$$

□

We now turn our attention to property (ii) of Δ_{2+}. Note that (ii) is trivially true because of the requirement that every function satisfying Δ_{2+} is twice differentiable, which can be easily perturbed. But even more is true.

Lemma 2.5. *Let M be an Orlicz function which satisfies condition Δ_{2+} and let $e > 0$ be given. Then there exists a twice differentiable Orlicz function M_1 such that*

$$(1-\varepsilon)M(u) \leq M_1(u) \leq (1+\varepsilon)M(u)$$

for all $u \in \mathbb{R}$ and which does not satisfy condition Δ_{2+}.

Sketch of proof. This fact is not of a particular importance for our paper so we do not provide a detailed proof but only sketch the idea.

Let M be an Orlicz function which satisfies (2.1) and (2.4). For any $n \in \mathbb{N}$ put

$$\begin{aligned} \alpha(n) &= M'(n + \frac{1}{2} + \frac{1}{2^{2n}}) - M'(n + \frac{1}{2} - \frac{1}{2^{2n}}) \\ \beta(n) &= \frac{\alpha(n)}{2^n} \frac{n + \frac{1}{2} - \frac{1}{2^{2n}}}{M'(n + \frac{1}{2} + \frac{1}{2^{2n}})}. \end{aligned}$$

Then there exists $n_0 \in \mathbb{N}$ so that for all $n \geq n_0$

$$\beta(n) < \frac{1}{2}\frac{1}{2^n}.$$

Indeed, by Mean Value Theorem, for each n there exists $c(n) \in (n+\frac{1}{2}-\frac{1}{2^{2n}}, n+\frac{1}{2}-\frac{1}{2^{2n}})$ so that $M''(c(n)) = \alpha(n)\cdot 2^n$. Then, by (2.4),

$$\beta > \frac{c(n)M''(c(n))}{M'(c(n))} = 2^n\alpha(n)\frac{c(n)}{M'(c(n))} \geq \beta(n)\cdot 2^{2n}.$$

Also this calculation implies that

$$\begin{aligned}\alpha(n) &< \frac{\beta M'(c(n))}{2^n c(n)} \\ &\text{by (2.1)} \\ &< \frac{\beta\alpha M(c(n))}{2^n (c(n))^2} \\ &\text{by (2.2)} \\ &\leq \frac{\beta\alpha C (c(n))^\alpha}{2^n (c(n))^2} \\ &\leq \frac{\beta\alpha C (n+1)^\alpha}{2^n n^2}.\end{aligned}$$

Thus $\alpha(n) \longrightarrow 0$ as $n \longrightarrow \infty$ and, for a given $\varepsilon > 0$ we can choose $n_1 \geq n_0$ so that for all $n \geq n_1$

$$\alpha(n) < \varepsilon.$$

For $n \geq n_1$ we "adjust" M' on $(n, n+1)$ by setting (see the graph in Figure 2):

$$\tilde{M}'(x) = \begin{cases} M'(n+\frac{1}{2}-\frac{1}{2^{2n}}) & \text{if } x \in (n+\frac{1}{2}-\frac{1}{2^{2n}}, n+\frac{1}{2}-\frac{\beta(n)}{2}) \text{ for some } n \geq n_1, \\ M'(n+\frac{1}{2}+\frac{1}{2^{2n}}) & \text{if } x \in (n+\frac{1}{2}+\frac{\beta(n)}{2}, n+\frac{1}{2}+\frac{1}{2^{2n}}) \text{ for some } n \geq n_1, \\ M'(n+\frac{1}{2}-\frac{1}{2^{2n}}) + (x-(n+\frac{1}{2}-\frac{\beta(n)}{2}))\cdot\frac{\alpha(n)}{\beta(n)} & \\ & \text{if } x \in (n+\frac{1}{2}-\frac{\beta(n)}{2}, n+\frac{1}{2}+\frac{\beta(n)}{2}) \text{ for some } n \geq n_1, \\ M'(x) & \text{otherwise.} \end{cases}$$



Figure 2.

$\tilde{M}'$ is not differentiable at countable number of points (four on each interval $(n, n+1)$ when $n \geq n_1$), but it can be further adjusted to produce M_1' so that M_1' is differentiable everywhere, $|M_1'(x) - M'(x)| < \varepsilon$ for all $x \geq 0$ and

$$M_1'(n+\frac{1}{2}) = \tilde{M}'(n+\frac{1}{2}),$$

$$M_1''(n+\frac{1}{2}) = \tilde{M}''(n+\frac{1}{2}) = \frac{\alpha(n)}{\beta(n)}.$$

We leave up to an interested reader the detailed formula for M_1 and the check that M_1 is $(1+\varepsilon)-$equivalent with M.

For M_1 we have for $n \geq n_1$

$$\frac{(n+\frac{1}{2})M_1''(n+\frac{1}{2})}{M_1'(n+\frac{1}{2})} = \frac{(n+\frac{1}{2})\alpha(n)M'(n+\frac{1}{2}+\frac{1}{2^{2n}})}{M_1'(n+\frac{1}{2})\alpha(n)(n+\frac{1}{2}-\frac{1}{2^{2n}})} \geq 2^n.$$

Hence M_1 does not satisfy condition Δ_{2+}. $\square$

In the case when M satisfies the Δ_2-condition and $\|f\|_M < \infty$ we have

$$\int_\Omega M\left(\frac{|f(u)|}{\|f\|_M}\right) d\mu = 1.$$

When M satisfies condition Δ_2 then the dual space of L_M is also an Orlicz space, which is determined by an Orlicz function M^* called *a complementary Orlicz function to* M which is defined by

$$M^*(v) = \int_0^v q(s)\ ds$$

for $v \geq 0$, where q denotes the right inverse of M' – the right derivative of M.

It is well-known (see [**KrR**]) that M^* does not have to satisfy condition Δ_2 whenever M does and also vice-versa: M^* may satisfy condition Δ_2 when M does not. Notice, however, that when M is twice differentiable, M'' is continuous and $M''(t) > 0$ for all $t > 0$ then the same is true for M^*, i.e. M^* is twice differentiable, $M^{*''}$ is continuous and $M^{*''}(t) > 0$ for all $t > 0$. Moreover, in this situation, if $\lim_{t\to 0} M''(t) = \infty$ then $\lim_{t\to 0} M^{*''}(t) = 0$.

For any $f \in L_M$ we will denote by f^N the *norming functional* for f i.e. the functional such that $\|f\|_{L_M} = \|f^N\|_{L^*_M}$ and $f^N(f) = \|f\|^2_{L_M}$.

We will need the following simple fact:

LEMMA 2.6. *Let M be a differentiable Orlicz function which satisfies condition Δ_2, A, B be disjoint subsets of* $[0,1]$ *and* $H = \operatorname{span}\{\chi_A, \chi_B\}$. *Suppose that* $T : L_M \xrightarrow{into} L_M$ *is an isometry.*

Then $(TH)^* = \{(Th)^N\ :\ h \in H\} = \operatorname{span}\{(T\chi_A)^N, (T\chi_B)^N\}$ *and the map* $S : \operatorname{span}\{\chi_A^N, \chi_B^N\} \longrightarrow \operatorname{span}\{(T\chi_A)^N, (T\chi_B)^N\}$ *defined by*

$$S\chi_A^N = (T\chi_A)^N, \qquad S\chi_B^N = (T\chi_B)^N$$

is an isometry.

PROOF. Since M is differentiable and satisfies condition Δ_2, by [**GH**], L_M is smooth. Since L_M is a symmetric function space, for every $A \subset [0,1]$, the norming functional $\chi_A^N = a\chi_A$ for an appropriate constant a. Indeed, if σ is a homeomorphism of $[0,1]$ with $\sigma(A) = A$, then $Tf = f \circ \sigma$ is a surjective isometry of L_M such that $T\chi_A = \chi_A$ and $(T^{-1})^*\chi_A^N = (T\chi_A)^N = \chi_A$. Thus

$$\chi_A^N = \chi_A^N \circ \sigma^{-1}$$

for any homeomorphism of $[0,1]$ with $\sigma(A) = A$. Hence $\chi_A^N\big|_A = const.$

Similarly, for any disjoint subsets $A, B \subset [0,1]$, by considering homeomorphisms of $[0,1]$ with $\sigma(A) = A$ and $\sigma(B) = B$ and the isometries of L_M that they induce, we conclude that for every $h \in H = \operatorname{span}\{\chi_A, \chi_B\}$, the norming functional for h is a linear combination of norming functionals χ_A^N, χ_B^N and $H^* = \operatorname{span}\{\chi_A^N, \chi_B^N\}$.

Let $V = T^{-1} : TL_M \longrightarrow L_M$. Since L_M is smooth for all $f \in L_M$ we have:

$$(Tf)^N = V^*(f^N).$$

Thus for every $h \in H$:

$$(Th)^N = V^*(h^N) \in \operatorname{span}\{V^*(\chi_A^N), V^*(\chi_B^N)\} = \operatorname{span}\{(T\chi_A)^N, (T\chi_B)^N\}$$

and

$$(TH)^* = \operatorname{span}\{(T\chi_A)^N, (T\chi_B)^N\}.$$

Further $V^* : H^* \longrightarrow (TH)^*$ is an isometry between subspaces of L_{M^*} with

$$V^*\chi_A^N = (T\chi_A)^N \quad , \quad V^*\chi_B^N = (T\chi_B)^N.$$

□

We finish this section with a lemma about differentiability of the function

$$F(\alpha, \eta) = \int M\left(\frac{|f + \alpha g|}{\eta}\right) d\mu(t) - 1, \tag{2.6}$$

where $\alpha \in \mathbb{R}$, $\eta > 0$ and f, g are given functions of norm 1 from L_M. We will need this lemma to describe the differential behaviour of $N(\alpha) = \|f + \alpha g\|_M$. Lemma 2.7 is inspired by and generalizes [**Kol**, Lemma 1].

Lemma 2.7. *Let M be an Orlicz function satisfying condition Δ_{2+} and such that M'' is continuous. Suppose that $f, g \in L_M$ with $\|f\| = \|g\| = 1$. For $\alpha \in \mathbb{R}$, $\eta > 0$, consider the function F defined by* (2.6)
Then

(a) $\frac{\partial^2 F}{\partial \alpha^2}(\alpha, \eta)$ *is a continuous function with respect to η, when $\eta \in (0, 1)$ and*

$$\frac{\partial^2 F}{\partial \alpha^2}(\alpha, 1) = \int M''(|f + \alpha g|) g^2 \, d\mu(t)$$

for a.e. $\alpha \in \mathbb{R}$,

(b) $\frac{\partial F}{\partial \eta}(\alpha, \eta)$ *is a continuous function with respect to both variables and*

$$0 < \left|\frac{\partial F}{\partial \eta}(0, 1)\right| < \infty,$$

(c) $\frac{\partial^2 F}{\partial \alpha \partial \eta}(\alpha, \eta)$ *and* $\frac{\partial^2 F}{\partial \eta^2}(\alpha, \eta)$ *are continuous functions with respect to both variables and*

$$\left|\frac{\partial^2 F}{\partial \alpha \partial \eta}(0, 1)\right| < \infty, \qquad \left|\frac{\partial^2 F}{\partial \eta^2}(0, 1)\right| < \infty.$$

Proof. The proofs of parts $(a), (b), (c)$ are very similar to each other and essentially consist of an application of Fubini's and Lebesgue's theorems.

We will need the following auxiliary functions, for $\alpha \in \mathbb{R}$, $\eta > 0$:

$$\begin{aligned} h_\eta(\alpha) &= \int M'\left(\frac{|f + \alpha g|}{\eta}\right) \operatorname{sgn}(f + \alpha g) \frac{g}{\eta} \, d\mu(t) \ , \\ s_\eta(\alpha) &= \int M''\left(\frac{|f + \alpha g|}{\eta}\right) \frac{g^2}{\eta^2} \, d\mu(t) \ , \\ w_\alpha(\eta) &= \int M'\left(\frac{|f + \alpha g|}{\eta}\right) \left(-\frac{|f + \alpha g|}{\eta^2}\right) d\mu(t) \ , \end{aligned}$$

$$\begin{aligned} v(\alpha,\eta) &= -\int M''\left(\frac{|f+\alpha g|}{\eta}\right)\frac{(f+\alpha g)g}{\eta^3}\,d\mu(t) \\ &\quad -\int M'\left(\frac{|f+\alpha g|}{\eta}\right)\operatorname{sgn}(f+\alpha g)\frac{g}{\eta^2}\,d\mu(t), \\ z_\alpha(\eta) &= \int M''\left(\frac{|f+\alpha g|}{\eta}\right)\frac{|f+\alpha g|^2}{\eta^4}\,d\mu(t) \\ &\quad +\int M'\left(\frac{|f+\alpha g|}{\eta}\right)\frac{2|f+\alpha g|}{\eta^3}\,d\mu(t) \end{aligned}$$

First notice that, since M satisfies condition (2.4), there exists $\beta > 0$ so that for all $\alpha \in \mathbb{R}$

$$M''\left(\frac{|f+\alpha g|}{\eta}\right)|f+\alpha g| \le \beta M'\left(\frac{|f+\alpha g|}{\eta}\right).$$

By [**KrR**, Lemma 9.1, p.73], for any $\alpha \in \mathbb{R}$ and $\eta > 0$

$$M'\left(\frac{|f+\alpha g|}{\eta}\right) \in L_{M^*}\ ,$$

where M^* is the complementary Orlicz function to M. Thus, by the generalized Hölder's inequality ([**KrR**, Theorem 9.3, p.74]):

$$|h_\eta(\alpha)| < \infty\ , |w_\alpha(\eta)| < \infty\ , |v(\alpha,\eta)| < \infty\ , |z_\alpha(\eta)| < \infty, \tag{2.7}$$

for all $\alpha \in \mathbb{R}$, $\eta > 0$.

Notice that for any $\beta, u, v \in \mathbb{R}$, $\eta > 0$ we have:

$$\int_0^\beta M''\left(\frac{|u+\alpha v|}{\eta}\right)\frac{v^2}{\eta^2}\,d\alpha = M'\left(\frac{|u+\alpha v|}{\eta}\right)\operatorname{sgn}(u+\alpha v)\frac{v}{\eta}\Big|_0^\beta$$

So, by the Fubini theorem

$$\int_0^\beta s_\eta(\alpha)\,d\alpha = h_\eta(\beta) - h_\eta(0)\ .$$

Thus s_η is absolutely integrable on $[0,\beta]$ with respect to α, $|s_\eta| < \infty$ for almost all α and h_η is a primitive for s_η. By Lebesgue's theorem, for each $\eta > 0$,

$$\frac{d}{d\alpha}h_\eta(\alpha) = s_\eta(\alpha)$$

for almost all $\alpha \in \mathbb{R}$.

Similarly, for each $\eta > 0$, $\int_0^\beta h_\eta(\alpha)\,d\alpha = F(\alpha,\eta)|_0^\beta$ and

$$\frac{\partial F}{\partial \alpha}(\alpha,\eta) = h_\eta(\alpha)$$

for almost all $\alpha \in \mathbb{R}$. Thus for $\eta = 1$

$$s_1(\alpha) = \frac{d}{d\alpha}h_1(\alpha) = \frac{\partial}{\partial\alpha}\frac{\partial F}{\partial\alpha}(\alpha,\eta)$$

and we get the formula in part (a).

The continuity of $\frac{\partial^2 F}{\partial\alpha^2}(\alpha,\eta)$ follows from the fact that M'' is continuous (see e.g. [**C**, Sections IV.2, IV.4]).

To prove (b) we use the same argument after we notice that for each $\alpha \in \mathbb{R}$ and for any $\varepsilon, \zeta > 0$:

$$\int_\varepsilon^\zeta M'\left(\frac{|u+\alpha v|}{\eta}\right)\left(-\frac{|u+\alpha v|}{\eta^2}\right)d\eta = M\left(\frac{|u+\alpha v|}{\eta}\right)\Big|_\varepsilon^\zeta.$$

Thus, as above, $\frac{\partial F}{\partial \eta}(\alpha,\eta) = w_\alpha(\eta)$ for almost all $\eta > 0$ and the continuity of $\frac{\partial F}{\partial \eta}(\alpha,\eta)$ follows from the continuity of M' [**C**].

Since $w_\alpha(\eta) < 0$ for all $\eta > 0$, we get $\frac{\partial F}{\partial \eta}(\alpha,1) \neq 0$.

Similarly, to get (c) we repeat the argument from above since we have for each $\eta > 0$, and $\varepsilon, \zeta > 0$

$$\int_\varepsilon^\zeta \left(-M''\left(\frac{|u+\alpha v|}{\eta}\right)\frac{(u+\alpha v)v}{\eta^3} - M'\left(\frac{|u+\alpha v|}{\eta}\right)\operatorname{sgn}(u+\alpha v)\frac{v}{\eta^2}\right)d\alpha$$

$$= \left(-M'\left(\frac{|u+\alpha v|}{\eta}\right)\frac{|u+\alpha v|}{\eta^2}\right)\Big|_{\alpha=\varepsilon}^{\alpha=\zeta}$$

Thus

$$v(\alpha,\eta) = \frac{\partial}{\partial\alpha}w_\alpha(\eta) = \frac{\partial^2 F}{\partial\alpha\partial\eta}(\alpha,\eta)$$

for almost all $\alpha \in \mathbb{R}, \eta > 0$.

Finally, for each $\eta > 0$, and $\varepsilon, \zeta > 0$

$$\int_\varepsilon^\zeta \left(M''\left(\frac{|u+\alpha v|}{\eta}\right)\frac{|u+\alpha v|^2}{\eta^4} + M'\left(\frac{|u+\alpha v|}{\eta}\right)\frac{2|u+\alpha v|}{\eta^3}\right)d\eta$$

$$= \left(-M'\left(\frac{|u+\alpha v|}{\eta}\right)\frac{|u+\alpha v|}{\eta^2}\right)\Big|_{\eta=\varepsilon}^{\eta=\zeta}$$

Thus

$$z_\alpha(\eta) = \frac{\partial}{\partial\eta}w_\alpha(\eta) = \frac{\partial^2 F}{\partial\eta^2}(\alpha,\eta)$$

for almost all $\alpha \in \mathbb{R}, \eta > 0$.

Continuity of both $v(\alpha,\eta)$ and $z_\alpha(\eta)$ with respect to α and η is again a consequence of continuity of M' and M''. The final statement follows from (2.7). ☐

3. The case of $M''(0) = 0$

We first study the case of Orlicz spaces L_M analogous to $L_p, p > 2$ in a sense that $M''(0) = 0$. We obtain the following partial description of functions with disjoint supports.

PROPOSITION 3.1. *Assume that M is an Orlicz function which satisfies condition Δ_{2+} and such that M'' is a continuous function with $M''(0) = 0$ and $M''(t) > 0$ for all $t > 0$. Let $f, g \in L_M$ and $N(\alpha) = \|f + \alpha g\|_M$.*

Then

(a) *If f, g have disjoint supports and g is bounded then $N'(0) = 0$ and $N''(\alpha) \longrightarrow 0$ as $\alpha \longrightarrow 0$ along a subset of $[0,1]$ of full measure.*

(b) *If $N'(0) = 0$ and $N''(\alpha) \longrightarrow 0$ as $\alpha \longrightarrow 0$ along a subset of $[0,1]$ of full measure then f, g have disjoint supports.*

PROOF. First note that since M is differentiable and satisfies condition Δ_2 thus, by [**GH**], L_M is smooth and the function $N(\alpha)$ is differentiable. Since $N(\alpha)$ is a convex function of α also the second derivative $N''(\alpha)$ exists a.e.

Notice that if $f, g \in L_M$ are disjointly supported then N clearly has a minimum at 0, so $N'(0) = 0$. Thus in the following we will work under the assumption that $N'(0) = 0$.

Assume, without loss of generality that $\|f\|_M = \|g\|_M = 1$, and let $F(\alpha, \eta)$ be defined as in Lemma 2.7. Since M satisfies Δ_2-condition we have $F(\alpha, N(\alpha)) = 0$ for all $\alpha \in \mathbb{R}$. Therefore $\frac{d}{d\alpha}(F(\alpha, N(\alpha))) = 0$ for all α. Hence

$$0 = \frac{d}{d\alpha}(F(\alpha, N(\alpha))) = \frac{\partial F}{\partial \alpha}(\alpha, N(\alpha)) + \frac{\partial F}{\partial \eta}(\alpha, N(\alpha))N'(\alpha) \tag{3.1}$$

and by taking the derivative again we get:

$$\begin{aligned} -\frac{\partial^2 F}{\partial \alpha^2}(\alpha, N(\alpha)) = {} & \frac{\partial^2 F}{\partial \alpha \partial \eta}(\alpha, N(\alpha))N'(\alpha) \\ & + \left[\frac{\partial^2 F}{\partial \alpha \partial \eta}(\alpha, N(\alpha)) + \frac{\partial^2 F}{\partial \eta^2}(\alpha, N(\alpha))N'(\alpha)\right] N'(\alpha) \\ & + \frac{\partial F}{\partial \eta}(\alpha, N(\alpha))N''(\alpha) \end{aligned} \tag{3.2}$$

Let A be the set so that for $\alpha \in A$, $N'(\alpha)$ exists and $\frac{\partial^2 F}{\partial \alpha^2}(\alpha, 1)$ is given by the formula from Lemma 2.7(a). By Lemma 2.7(c) there exists a sequence $(\alpha_n) \subset A, \alpha_n \to 0$ so that $\frac{\partial^2 F}{\partial \alpha \partial \eta}(\alpha_n, N(\alpha_n))$ and $\frac{\partial^2 F}{\partial \eta^2}(\alpha_n, N(\alpha_n))$ are bounded for all $n \in \mathbb{N}$. Since $N'(\alpha_n) \to 0$ and since, by Lemma 2.7(b), $\frac{\partial F}{\partial \eta}(\alpha_n, N(\alpha_n)) \neq 0$ for n large enough, we get that

$$\lim_{n\to\infty} \frac{\partial^2 F}{\partial \alpha^2}(\alpha_n, N(\alpha_n)) = 0 \Longleftrightarrow \lim_{n\to\infty} N''(\alpha_n) = 0 \ .$$

Since $\frac{\partial^2 F}{\partial \alpha^2}(\alpha, \eta)$ is a continuous function with respect to both α and η we conclude that

$$\lim_{n\to\infty} \frac{\partial^2 F}{\partial \alpha^2}(\alpha_n, 1) = 0 \Longleftrightarrow \lim_{n\to\infty} N''(\alpha_n) = 0 \ .$$

Since $\alpha_n \in A$, we obtain, by Lemma 2.7(a)

$$\frac{\partial^2 F}{\partial \alpha^2}(\alpha_n, 1) = \int M''(|f + \alpha_n g|)g^2 d\mu(t) \ . \tag{3.3}$$

If $\operatorname{supp} f \cap \operatorname{supp} g = \emptyset$ then (3.3) becomes

$$\frac{\partial^2 F}{\partial \alpha^2}(\alpha_n, 1) = \int M''(|\alpha_n g|)g^2 d\mu(t)$$

and since $|g|$ is bounded and $M''(t) \to 0$ as $t \to 0$ we conclude that $\frac{\partial^2 F}{\partial \alpha^2}(\alpha_n, 1) \longrightarrow 0$ as $n \to \infty$ and therefore $\lim_{n\to\infty} N''(\alpha_n) = 0$.

If $\operatorname{supp} f \cap \operatorname{supp} g \neq \emptyset$ let $B \subset \operatorname{supp} f \cap \operatorname{supp} g$ and $b_1, b_2, b_3, b_4 > 0$ be such that $\mu(B) > 0$, $0 < b_1 < |f(t)| < b_2$, $0 < b_3 < |g(t)| < b_4$ for all $t \in B$. Now let $n_0 \in \mathbb{N}$ be such that for all $n > n_0$ and all $t \in B$:

$$M''(|f(t) + \alpha_n g(t)|)) > \frac{1}{2} M''(b_1) \ .$$

Then since $M''(|f(t)+\alpha_n g(t)|)g^2(t) \geq 0$ for a.e. t, we get for $n > n_0$:

$$\int M''(|f+\alpha_n g|)g^2 d\mu(t) \geq \int_B M''(|f+\alpha_n g|)g^2 d\mu(t) \geq \frac{1}{2}\mu(B)M''(b_1)b_3^2 > 0 \ .$$

Hence $\lim_{n\to\infty} \frac{\partial^2 F}{\partial \alpha^2}(\alpha_n, 1) \neq 0$ and therefore $\lim_{n\to\infty} N''(\alpha_n) \neq 0$. ☐

The above partial characterization of disjointness allows us to immediately conclude that isometries from subspaces of L_M which contain enough disjointly supported bounded functions into L_M have to preserve disjointness.

THEOREM 3.2. *Assume that M is an Orlicz function which satisfies condition Δ_{2+} and such that M'' is a continuous function with $M''(0) = 0$. Let $T : L_M \overset{into}{\longrightarrow} L_M$ be an isometry. Then T preserves disjointness.*

4. The case of $M''(0) = \infty$

In this section we study Orlicz spaces L_M analogous to $L_p, 1 < p < 2$. In this case we do not find any characterizations of disjointness. Instead we give conditions which help us to determine when support of f is contained in the support of g. Our conditions do not provide a full characterization of containment of supports but they are sufficient to determine that isometries have to preserve the containment of supports.

PROPOSITION 4.1. *Assume that M is an Orlicz function which satisfies Δ_{2+} condition and such that M'' is a continuous function on $(0,\infty)$ with $\lim_{t\to 0} M''(t) = \infty$. Let $f, g \in L_M$ with $\|f\| = \|g\| = 1$.*

Then

(a) $\mu(\operatorname{supp} g \setminus \operatorname{supp} f) > 0 \Rightarrow \lim_{\alpha\to 0} N''(\alpha) = \infty$

(b) *If f, g are simple then*

$$\mu(\operatorname{supp} g \setminus \operatorname{supp} f) = 0 \Rightarrow \lim_{\alpha\to 0} N''(\alpha) \neq \infty \ .$$

PROOF. Similarly as in the proof of Proposition 3.1 we see that equation (3.2) is valid i.e.

$$\begin{aligned} -\frac{\partial^2 F}{\partial \alpha^2}(\alpha, N(\alpha)) &= \frac{\partial^2 F}{\partial \alpha \partial \eta}(\alpha, N(\alpha))N'(\alpha) \\ &+ \left[\frac{\partial^2 F}{\partial \alpha \partial \eta}(\alpha, N(\alpha)) + \frac{\partial^2 F}{\partial \eta^2}(\alpha, N(\alpha))N'(\alpha)\right] N'(\alpha) \\ &+ \frac{\partial F}{\partial \eta}(\alpha, N(\alpha))N''(\alpha) \end{aligned} \tag{4.1}$$

Let A be the set so that for $\alpha \in A$, $N''(\alpha)$ exists and $\frac{\partial^2 F}{\partial \alpha^2}(\alpha, 1)$ is given by the formula from Lemma 2.7(a). By Lemma 2.7(c) there exists a sequence $(\alpha_n) \subset A, \alpha_n \to 0$ so that $\frac{\partial^2 F}{\partial \eta \partial \alpha}(\alpha_n, N(\alpha_n))$ and $\frac{\partial^2 F}{\partial \eta^2}(\alpha_n, N(\alpha_n))$ are bounded for all $n \in \mathbb{N}$. $N'(\alpha)$ is a continuous function of α and $|N'(0)| < \infty$ by (3.1) and Lemma 2.7(b). Thus $\frac{\partial^2 F}{\partial \eta \partial \alpha}(\alpha_n, N(\alpha_n))N'(\alpha_n)$ and $\frac{\partial^2 F}{\partial \eta^2}(\alpha_n, N(\alpha_n))N'(\alpha_n)$ are bounded for all $n \in \mathbb{N}$. Therefore, since $\frac{\partial F}{\partial \eta}(\alpha_n, N(\alpha_n))$ is bounded for all n by Lemma 2.7(b), we get from (4.1) that

$$\lim_{n\to\infty} \frac{\partial^2 F}{\partial \alpha^2}(\alpha_n, N(\alpha_n)) = \infty \iff \lim_{n\to\infty} N''(\alpha_n) = \infty \ .$$

Since $\frac{\partial^2 F}{\partial\alpha^2}(\alpha_n,\eta)$ is a continuous function with respect to η we conclude that

$$\lim_{n\to\infty}\frac{\partial^2 F}{\partial\alpha^2}(\alpha_n,1)=\infty \iff \lim_{n\to\infty}N''(\alpha_n)=\infty \ .$$

Since $\alpha_n\in A$, we obtain by Lemma 2.7(a)

$$\frac{\partial^2 F}{\partial\alpha^2}(\alpha_n,1)=\int_{\operatorname{supp} g}M''(|f+\alpha_n g|)g^2d\mu(t)$$

For (a) assume that $\mu(\operatorname{supp} g\setminus\operatorname{supp} f)>0$ and let $S\subset\operatorname{supp} g\setminus\operatorname{supp} f$ be such that $\mu(S)>0, \inf\{|g(t)| : t\in S\}=s_1>0$ and $\sup\{|g(t)| : t\in S\}=s_2<\infty$. Denote

$$m_n=\inf\{M''(|\alpha_n s|) \ : \ s\in[s_1,s_2]\}.$$

Since $\alpha_n\to 0$ as $n\to\infty$ and $M''(t)\to\infty$ as $t\to 0$, we get $\lim_{n\to\infty}m_n=\infty$. Thus we have

$$\begin{aligned}\frac{\partial^2 F}{\partial\alpha^2}(\alpha_n,1) &\geq \int_S M''(|\alpha_n g|)g^2d\mu(t)\\ &\geq \mu(S)m_n\cdot s_1^2\underset{n\to\infty}{\longrightarrow}\infty.\end{aligned}$$

Thus $\lim_{n\to\infty}N''(\alpha_n)=\infty$ and therefore $\lim_{\alpha\to 0}N''(\alpha)=\infty$.

For (b) assume that f,g are simple and $\mu(\operatorname{supp} g\setminus\operatorname{supp} f)=0$. Set $m_f=\inf\{f(t) : t\in\operatorname{supp} g\}$. Since f is a simple function $m_f>0$ and since g is bounded there exists $\alpha_0>0$ so that $|f(t)+\alpha g(t)|>\frac{1}{2}m_f$ for all t and all α with $|\alpha|<\alpha_0$. Let $\delta_0=\sup\{M''(t) : \frac{1}{2}m_f\leq t\leq\|f\|_\infty+\|g\|_\infty\}$. By continuity of M'' on $(0,\infty), \delta_0<\infty$. Thus for all n such that $|\alpha_n|<\alpha_0$ we have:

$$\int_{\operatorname{supp} g}M''(|f+\alpha_n g|)g^2d\mu(t)\leq\delta_0\cdot\|g\|_\infty^2<\infty$$

Hence $\lim_{n\to\infty}N''(\alpha_n)\neq\infty$ and therefore $\lim_{\alpha\to 0}N''(\alpha)\neq\infty$. □

Corollary 4.2. *Assume that M is an Orlicz function which satisfies Δ_{2+} condition and such that M'' is a continuous function on $(0,\infty)$ with $\lim_{t\to 0}M''(t)=\infty$. Let $T: L_M\overset{into}{\longrightarrow}L_M$ be an isometry and $f,g\in L_M$ be such that $\operatorname{supp} f=\operatorname{supp} g$ up to a set of measure zero. Then $\operatorname{supp} Tf=\operatorname{supp} Tg$ up to a set of measure zero.*

Proof. Without loss of generality we assume that f,g are simple and $\|f\|=\|g\|=1$. By Proposition 4.1(b) we see that $\lim_{\alpha\to 0}N''(\alpha)\neq\infty$. Since T is an isometry $N(\alpha)=\|Tf+\alpha Tg\|$ and thus by Proposition 4.1(a) $\mu(\operatorname{supp} Tg\setminus\operatorname{supp} Tf)=0$.

Exchanging the roles of f and g we symmetrically obtain $\mu(\operatorname{supp} Tf\setminus\operatorname{supp} Tg)=0$. □

After the author presented this paper at the Conference on Function Spaces at Southern Illinois University at Edwardsville, Abramovich and Kitover [**AK1**] showed that an isometry T between Banach function spaces satisfies condition from the conclusion of Corollary 4.2 if and only if $T^{-1}: TX\longrightarrow X$ is disjointness preserving. However they also showed that in general it is possible to construct operators satisfying the above condition but such that TX does not have non-trivial disjoint elements and, in particular, T is not disjointness preserving (cf. also [**AK2**]).

Below we show that such a situation cannot happen in the case of Orlicz spaces, i.e. every injective isometry preserving equality of supports does preserve disjointness.

THEOREM 4.3. *Assume that M is an Orlicz function which satisfies Δ_{2+} condition and such that M'' is a continuous function on $(0,\infty)$ and $\lim_{t\to 0} M''(t)=\infty$. Then every isometry $T: L_M \overset{into}{\longrightarrow} L_M$ preserves disjointness.*

PROOF. By Corollary 4.2 it is enough to show that for any two disjoint sets $A, B, \subset [0,1]$ we have $\mu(\operatorname{supp} T\chi_A \cap \operatorname{supp} T\chi_B) = 0$.

The proof of this fact is very short if we assume in addition that $M''(t) > 0$ for all $t > 0$ and that the complementary function M^* satisfies Δ_{2+} condition. We present this simple duality argument first, and then we show a longer proof which does not require any additional assumptions.

Denote $H = \operatorname{span}\{\chi_A, \chi_B\}$. By Lemma 2.6 the map $S : \operatorname{span}\{\chi_A^N, \chi_B^N\} \longrightarrow \operatorname{span}\{(T\chi_A)^N, (T\chi_B)^N\}$ defined by

$$S\chi_A^N = (T\chi_A)^N, \qquad S\chi_B^N = (T\chi_B)^N$$

is an isometry.

Notice that for all v

$$(M^*)''(v) = \frac{1}{M''\big((M')^{-1}(v)\big)}.$$

So $(M^*)''$ is continuous, $(M^*)''(v) > 0$ for all $v > 0$ and

$$\lim_{v\to 0}(M^*)''(v) = \lim_{\omega\to 0}\frac{1}{M''(\omega)} = 0.$$

If M^* satisfies Δ_{2+}-condition we obtain by Proposition 3.1 that S preserves disjointness, since bounded functions are clearly dense in $\operatorname{span}\{\chi_A^N, \chi_B^N\} \subset L_{M^*}$. And since $\operatorname{supp} h^N = \operatorname{supp} h$ for all $h \in L_M$, we conclude that $\mu(\operatorname{supp} T\chi_A \cap \operatorname{supp} T\chi_B) = 0$ as desired.

If M^* does not satisfy Δ_{2+} condition or if $M''(t) \not> 0$ for all $t > 0$, we will apply a much longer, more direct approach relying on the fact that $(TH)^* = \operatorname{span}\{(T\chi_A)^N, (T\chi_B)^N\}$ is a subspace of L_{M^*}.

Denote $f = \chi_A$, $g = \chi_B$ and for any scalar α let $h = \chi_A + \alpha\chi_B$ and $Th = f + \alpha g$. By [**GH**] for almost every $t \in [0,1]$ we have:

$$\begin{aligned}
(Th)^N(t) &= C_\alpha M'\left(\frac{|(f+\alpha g)(t)|}{\|f+\alpha g\|}\right)\operatorname{sgn}((f+\alpha g)(t)) \ , \\
f^N(t) &= C_f M'\left(\frac{|f(t)|}{\|f\|}\right)\operatorname{sgn}(f(t)) \ , \\
g^N(t) &= C_g M'\left(\frac{|g(t)|}{\|g\|}\right)\operatorname{sgn}(g(t)) \ ,
\end{aligned}$$

where constants C_α, C_f, C_g do not depend on t.

Thus for each α there exist $\beta_1(\alpha), \beta_2(\alpha)$ so that

$$M'\left(\frac{|(f+\alpha g)(t)|}{\|f+\alpha g\|}\right)\operatorname{sgn}((f+\alpha g)(t)) = \beta_1(\alpha)M'\left(\frac{|f(t)|}{\|f\|}\right)\operatorname{sgn}(f(t)) + \beta_2(\alpha)M'\left(\frac{|g(t)|}{\|g\|}\right)\operatorname{sgn}(g(t)). \tag{4.2}$$

Since $A \cap B = \emptyset$, by Corollary 4.2,

$$\begin{aligned} \mu(\operatorname{supp} g \setminus \operatorname{supp} f) &> 0 \text{ and} \\ \mu(\operatorname{supp} f \setminus \operatorname{supp} g) &> 0 . \end{aligned}$$

Thus for $t \in \operatorname{supp} f \setminus \operatorname{supp} g$ equation (4.2) becomes

$$M'\left(\frac{|f(t)|}{N(\alpha)}\right)\operatorname{sgn}(f(t)) = \beta_1(\alpha)M'\left(\frac{|f(t)|}{\|f\|}\right)\operatorname{sgn}(f(t)) + 0. \tag{4.3}$$

Our next goal is to evaluate $\beta_1(\alpha)$. We follow a technique similar to the one in the proof of [**R**, Theorem 6.1].

Suppose that $|f|\Big|_{\operatorname{supp} f \setminus \operatorname{supp} g}$ is not constant, that is there exist $t_1, t_2 \in \operatorname{supp} f \setminus \operatorname{supp} g$ so that $|f(t_1)|/\|f\| \neq |f(t_2)|/\|f\|$. Denote $x_i = |f(t_i)|/\|f\|,\ i = 1, 2$. Then (4.3) becomes

$$\begin{aligned} M'\left(x_1 \cdot \frac{\|f\|}{N(\alpha)}\right) &= \beta_1(\alpha)M'(x_1) \\ M'\left(x_2 \cdot \frac{\|f\|}{N(\alpha)}\right) &= \beta_1(\alpha)M'(x_2) \end{aligned}$$

Thus

$$\frac{M'\left(x_1 \cdot \frac{\|f\|}{N(\alpha)}\right)}{M'(x_1)} = \frac{M'\left(x_2 \cdot \frac{\|f\|}{N(\alpha)}\right)}{M'(x_2)} \tag{4.4}$$

for all $\alpha \in \mathbb{R}$. Notice that $N([0,\infty)) = [\|f\|, \infty)$. Thus $(\|f\|/N(\alpha)) \in (0, 1]$.

Set

$$u = x_2\frac{\|f\|}{N(\alpha)}, \quad \beta = \frac{M'(x_1)}{M'(x_2)}, \quad \xi = \frac{x_1}{x_2} .$$

Then (4.4) becomes

$$M'(\xi u) = \beta M'(u)$$

for all $u \in [0, x_2]$.

Thus

$$M(\xi u) = \beta\xi M(u)$$

for all $u \in [0, x_2]$ and by [**R**, Lemma 6.2] there exists $p \geq 1$ and constants C_1, C_2 so that for all $u \leq x_2$

$$C_2 u^p \leq M(u) \leq C_1 u^p .$$

Moreover, if $M(u) \not\equiv Cu^p$ for $u \in [0, x_2]$ then there exists $\gamma > 0$ such that

$$M'(a\gamma^k) = M'(a)M'(\gamma^k) \tag{4.5}$$

for all $a > 0, k \in \mathbb{Z}$ and $\xi \in \{\gamma^k \in \mathbb{Z}\}$.

Thus if $|f|\Big|_{\operatorname{supp} f \setminus \operatorname{supp} g}$ is not constant and if $M(u) \not\equiv Cu^p$ for u near zero then there exists $x > 0$ and γ satisfying (4.5) so that $|f(t)|/\|f\| \in \{x\gamma^k : k \in \mathbb{Z}\}$ for all $t \in \operatorname{supp} f \setminus \operatorname{supp} g$. Hence (4.3) becomes

$$M'\left(x\gamma^k \frac{\|f\|}{N(\alpha)}\right) = \beta_1(\alpha)M'(x\gamma^k).$$

By (4.5) we get

$$M'(\gamma^k)M'\left(x\frac{\|f\|}{N(\alpha)}\right) = \beta_1(\alpha)M'(x)M'(\gamma^k)\ .$$

Hence

$$\beta_1(\alpha) = \frac{M'\left(x\frac{\|f\|}{N(\alpha)}\right)}{M'(x)}\ . \tag{4.6}$$

Clearly, if $|f|\Big|_{\operatorname{supp} f \setminus \operatorname{supp} g} \equiv \text{const.} = x$ then (4.3) becomes

$$M'\left(x\frac{\|f\|}{N(\alpha)}\right) = \beta_1(\alpha)M'(x)$$

and (4.6) holds.

Using similar technique we show that when $M(u) \not\equiv Cu^p$ for u near zero for any p, then there exists $y > 0$ so that

$$\beta_2(\alpha) = \frac{M'\left(y\frac{|\alpha|\ \|g\|}{N(\alpha)}\right)}{M'(y)}\ . \tag{4.7}$$

Thus for $t \in \operatorname{supp} f \cap \operatorname{supp} g$, (4.2) becomes:

$$\begin{aligned} M'\left(\frac{|f(t)+\alpha g(t)|}{N(\alpha)}\right)&\operatorname{sgn}(f(t)+\alpha g(t)) = \\ &\frac{1}{M'(x)}M'\left(x\cdot\frac{\|f\|}{N(\alpha)}\right)\cdot M'\left(\frac{|f(t)|}{\|f\|}\right)\operatorname{sgn}(f(t)) \\ &+\frac{1}{M'(y)}M'\left(y\cdot\frac{|\alpha|\ \|g\|}{N(\alpha)}\right)M'\left(\frac{|g(t)|}{\|g\|}\right)\operatorname{sgn}(g(t))\ . \end{aligned} \tag{4.8}$$

Fix $t \in \operatorname{supp} f \cap \operatorname{supp} g$ and let $\alpha_0 > 0$ be small enough so that $\operatorname{sgn}(f(t) + \alpha g(t)) = \operatorname{sgn}(f(t))$ for all $\alpha \in (0, \alpha_0)$. We differentiate (4.8) with respect to α when $\alpha \in (0, \alpha_0)$:

$$
\begin{aligned}
&M''\left(\frac{|f(t)+\alpha g(t)|}{N(\alpha)}\right)\operatorname{sgn}(f(t))\cdot\frac{\operatorname{sgn}(f(t))g(t)N(\alpha)-N'(\alpha)|f(t)+\alpha g(t)|}{(N(\alpha))^2}=\\
&\quad\frac{1}{M'(x)}M'\left(\frac{|f(t)|}{\|f\|}\right)\cdot\operatorname{sgn}(f(t))\cdot M''\left(x\cdot\frac{\|f\|}{N(\alpha)}\right)\cdot\frac{(-x\|f\|)}{(N(\alpha))^2}\\
&\quad+\frac{1}{M'(y)}M'\left(\frac{|g(t)|}{\|g\|}\right)\operatorname{sgn}(g(t))\cdot M''\left(y\cdot\frac{\alpha\|g\|}{N(\alpha)}\right)\cdot\frac{y\|g\|N(\alpha)-N'(\alpha)y\alpha\|g\|}{(N(\alpha))^2}
\end{aligned}
$$

When α approaches zero we obtain:

$$
\begin{aligned}
&M''\left(\frac{|f(t)|}{\|f\|}\right)\operatorname{sgn}(f(t))\cdot\frac{\operatorname{sgn}(f(t))g(t)\|f\|-0}{\|f\|^2}=\\
&\qquad\frac{1}{M'(x)}M'\left(\frac{|f(t)|}{\|f\|}\right)\operatorname{sgn}(f(t))M''(x)\cdot\left(\frac{-x\|f\|}{\|f\|^2}\right)\\
&\qquad+\frac{1}{M'(y)}M'\left(\frac{|g(t)|}{\|g\|}\right)\operatorname{sgn}(g(t))\cdot\frac{y\|g\|\cdot\|f\|-0}{\|f\|^2}\cdot\lim_{\alpha\to 0}M''\left(\alpha\frac{y\|g\|}{N(\alpha)}\right)
\end{aligned}
$$

which is a contradiction since

$$\lim_{\alpha\to 0}M''\left(\alpha\cdot\frac{y\|g\|}{N(\alpha)}\right)=\infty$$

and all other quantities in the above equation are finite and nonzero.

Hence $\mu(\operatorname{supp} f\cap\operatorname{supp} g)=0$. ☐

5. Final remarks

In this section we summarize the results that we obtained:

Corollary 5.1. *Suppose that M is an Orlicz function satisfying condition Δ_{2+} and such that M'' is continuous and either $\lim_{t\to 0}M''(t)=\infty$ or $M''(0)=0$ and $M''(t)>0$ for all $t>0$. Suppose further that $T: L_M \overset{into}{\longrightarrow} L_M$ is an isometry. Then there exist a Borel map $\sigma:[0,1]\longrightarrow[0,1]$ and a function $a:[0,1]\longrightarrow\mathbb{R}$ (or, if L_M is complex, $a:[0,1]\longrightarrow\mathbb{C}$) so that for every $f\in L_M$ and almost every $t\in[0,1]$:*

$$Tf(t)=a(t)f(\sigma(t)) \tag{5.1}$$

Moreover $|a(t)|=1$ a.e. unless there exist constants $C_1, C_2, t_0>0$ and $p, 1<p<\infty$,

so that for all $t\le t_0$:

$$C_1t^p\le M(t)\le C_2t^p.$$

If such constants exist, but $(M(t)/t^p)\not\equiv const.$ on any interval containing 0 then there exist $A,\gamma>0$ so that for a.e. t, $|a(t)|=A\cdot\gamma^{k(t)}$, where $k(t)\in\mathbb{Z}$.

Proof. It follows from Theorems 3.2 and 4.3 that T preserves disjointness. Abramovich [**A**] proved that this implies that T is a weighted composition operator i.e. T has form (5.1).

To prove the "moreover part" we will use the nonatomic version of [**R**, Theorem 6.1] which we remind below:

Theorem 5.2. *Let M be a continuous Orlicz function and let $f, g \in L_M$ be disjoint elements such that $f, g \not\equiv 0$ and $\{h^N : h \in \operatorname{span}\{f, g\}\} = \operatorname{span}\{f^N, g^N\} \subset L_M^*$. Then one of three possibilities holds:*

1. *there exists a scalar k_f so that $|f(t)| = k_f$ for almost all $t \in \operatorname{supp} f$; or*
2. *there exists $p, 1 \leq p < \infty$, $C \geq 0$ and $t_0 > 0$ so that $M(t) = Ct^p$ for all $t \leq t_0$; or*
3. *there exist $p, 1 \leq p < \infty$ and constants $C_1, C_2, t_0, \gamma, k_f \geq 0$ so that $C_2 t_p \leq M(t) \leq C_1 t^p$ for all $t \leq t_0$ amd such that for almost all $t \in \operatorname{supp} f$ there exists $k(t) \in \mathbb{Z}$ with*
$$|f(t)| = k_f \cdot \gamma^{k(t)}$$

Remark 5.3. [**R**, Theorem 6.1] is stated and proven for sequence spaces ℓ_M, but the nonatomic version requires only very minor routine adjustments, so we leave them to the interested reader.

Now we recall that by Lemma 2.6, for any disjoint subsets A, B of $[0, 1]$ we have $\{(Th)^N : h \in \operatorname{span}\{\chi_A, \chi_B\}\} = \operatorname{span}\{(T\chi_A)^N, (T\chi_B)^N\}$. But we know that $T\chi_A, T\chi_B$ are disjointly supported so the "moreover part" of Corollary 5.1 follows directly from Theorem 5.2 applied to $f = T\chi_A, g = T\chi_B$. $\square$

Remark 5.4. The statement of Corollary 5.1 leaves open the case when M'' is continuous and $M''(0) = c$ for some c, $0 < c < \infty$. The function $M_2(t) = t^2$ belongs to this case and of course L_{M_2} has non-disjointness preserving injective and surjective isometries.

If $M(t) \not\equiv t^2$ on $[0, 1]$ then it is known that all surjective isometries are disjointness preserving; however our differential technique does not seem to provide enough information about injective isometries in this case. We feel that the hardest case would be to distinguish behavior in L_2 from L_M where $M(t) = t^2$ for all $t \leq a < 1$ but a is close to 1.

Remark 5.5. Our results deal with injective isometries where domain is entire L_M(see also the remark before Theorem 3.2). It would be interesting to determine if isometries from subspaces of L_M into L_M have to be disjointness preserving, as it is the case in $L_p, p \neq 2$ (cf. [**Kol**]); (note that when $M(t) = t^2$ for all $t \leq a < 1$, where a is large enough, then L_M contains an isometric copy of ℓ_2^2 [**R**, Example 3], so clearly injective isometries from the subspace of L_M do not have to preserve disjointness in this case).

References

[A] Y. A. Abramovich. Multiplicative representation of disjointness preserving operators. *Indag. Math.*, 45:265–279, 1983.

[AK1] Y. A. Abramovich and A. K. Kitover. A characterization of operators preserving disjointness in terms of their inverse. preprint.

[AK2] Y. A. Abramovich and A. K. Kitover. A solution to a problem on invertible disjointness preserving operators. *Proc. Amer. Math. Soc.*, 126:1501–1505, 1998.

[Ban] S. Banach. *Theorie des operations lineaires.* Monografie Mat. 1, Polish Scientific Publishers, Warsaw, 1932.

[Ch] Shutao Chen. Geometry of Orlicz spaces. *Dissertationes Math.*, 356:204 pages, 1996.

[C] R. Courant. *Differential and Integral Calculus*, volume II. Interscience Publishers, Inc., New York, 1953.

[FJ] R. J. Fleming and J. E. Jamison. Isometries of Banach spaces – a survey. In *Analysis, geometry and groups: a Riemann legacy volume*, pages 52–123. Hadronic Press, Palm Harbor, FL, 1993.

[GH] R. Grząślewicz and H. Hudzik. Smooth points of Orlicz spaces equipped with Luxemburg norm. *Math. Nachr.*, 155:31–45, 1992.

[JKL] J. Jamison, A. Kamińska, and P. K. Lin. Isometries in Musielak–Orlicz spaces II. *Studia Math.*, 104:75–89, 1993.

[KR] N. J. Kalton and B. Randrianantoanina. Surjective isometries of rearrangement-invariant spaces. *Quart. J. Math. Oxford*, 45:301–327, 1994.

[Kam] A. Kamińska. Isometries of Orlicz spaces. In *Proceedings of the Orlicz Memorial Conference*. The University of Mississippi, Oxford, Mississippi, 1991.

[Kol] A. L. Koldobskiî̂. Isometries of $L_p(X; L_q)$ and equimeasurability. *Indiana Univ. Math. J.*, 40:677–705, 1991.

[KrR] M. A. Krasnosel'skii and Ya. B. Rutickii. *Convex functions and Orlicz spaces.* P. Noordhoff LTD., Groningen, The Netherlands, 1961.

[Lam] J. Lamperti. On the isometries of some function spaces. *Pacific J. Math.*, 8:459–466, 1958.

[LT] J. Lindenstrauss and L. Tzafriri. *Classical Banach spaces, Vol. 2, Function spaces.* Springer–Verlag, Berlin–Heidelberg–New York, 1979.

[L1] G. Lumer. Isometries of reflexive Orlicz spaces. *Bull. Amer. Math. Soc.*, 68:28–30, 1962.

[L2] G. Lumer. On the isometries of reflexive Orlicz spaces. *Ann. Inst. Fourier*, 13:99–109, 1963.

[R] B. Randrianantoanina. 1-complemented subspaces of spaces with 1-unconditional bases. *Canadian J. Math.*, 49:1242–1264, 1997.

[Z] M. G. Zaidenberg. Groups of isometries of Orlicz spaces. *Soviet Math. Dokl.*, 17:432–436, 1976.

Department of Mathematics and Statistics, Miami University, Oxford, OH 45056
E-mail address: `randrib@@muohio.edu`